LOYOLA MARYMOUNT UNIVERSITY STUDENT HEALTH SERVICES

ENTRANCE REQUIREMENTS

ALL INCOMING STUDENTS MUST COMPLETE THE FOLLOWING BY

January5th 2026

If you are unable to meet this deadline please upload your information as soon as possible

- 1. Review the Consent for Treatment Form.
- 2. Review the Notice of Privacy Practices.
- 3. Review the Telehealth Consent Form.
- 4. Provide documentation of two measles/mumps/rubella (MMR) immunization or proof of immunity (blood titer).
- 5. Provide documentation of immunization for Meningococcal ACWY or Meningococcal B
- 6. Complete Tuberculosis (TB) Screening and/or testing.

Go to your MyHealth portal located at

(MyHealth.lmu.edu) login and click on the <u>MEDICAL</u> CLEARANCES Tab

located on the left hand side of your screen.

• Complete all items in your Medical Clearances section.

Please Note:

• Tuberculosis (TB) Screening and/or testing

You must submit a TB skin or blood test done within the past 12 months, if your country of origin is designated by the World Health Organization (WHO) as a country with a high incidence of tuberculosis OR you answered "yes" to any of the questions listed in the TB Risk Screening Questionnaire in the Medical Clearances section.

 If your TB skin or blood test is positive, you must also submit a chest x-ray report as proof you do not have active TB.

CONTACT STUDENT HEALTH SERVICES

Phone: 310.338.2881

Fax: 310.338.4417

Email:

lmuhealth@lmu.edu

Office:

Burns Recreation Center Ground-floor level, North

Mailing address:

1 LMU Drive, MS 8455, Los Angeles CA 90045

Questions?

Please review the Entrance Requirements FAQ's on our web page: Imu.edu/health

Allow 5 business days for SHS to process your forms.



HIGH RISK FACTORS FOR ACQUIRING TB ARE AS FOLLOW:

- · Close contact with persons known or suspected to have active TB disease
- Persons born in one of the countries listed below that have a high incidence of active TB disease
- Frequent or prolonged visits to one or more of the countries listed below with a high prevalence of TB disease
- A resident and/or employee of high-risk congregate settings (e.g., correctional facilities, long-term care facilities, and homeless shelters)
- A volunteer or health-care worker who served clients who are at increased risk for active TB disease
- A member of any of the following groups that may have an increased incidence of latent M. tuberculosis infection or active TB disease medically underserved, low-income, or abusing drugs or alcohol

COUNTRIES WITH HIGH INCIDENCE OF TUBERCULOSIS PER WORLD HEALTH ORGANIZATION

Afghanistan China, Hong Kong SAR Solomon Islands Guyana Morocco Algeria China, Macao SAR Haiti Mozambique Somalia Angola Colombia Honduras Myanmar South Africa Anguilla Comoros India Namibia South Sudan Argentina Congo Indonesia Nauru Sri Lanka Armenia Cote d'Ivoire Sudan Iraq Nepal Azerbaijan Democratic People's Kazakhstan Nicaragua Suriname Bangladesh Republic of Korea Kenya Niger Tajikistan Nigeria Belarus Democratic Republic Kiribati Tanzania (United Belize of the Congo Niue Republic of) Kyrgyzstan Benin Diibouti Lao People's Northern Mariana islands Thailand Bhutan Dominican Republic Democratic Republic Pakistan Timor-Leste Bolivia(Plurinational) Ecuador Lesotho Palau Togo State of) El Salvador Panama Trinidad & Liberia Bosnia and Equatorial Guinea Libya Papua New Guinea Tobago Herzegovina Eritrea Lithuania Paraguay Tunisia Turkmenistan Botswana eSwatini Madagascar Peru Brazil Ethiopia Malawi Philippines Tuvalu Uganda Brunei Darussalam Fiji Malaysia Qatar Maldives Republic of Korea Ukraine Burkina Faso Gabon Burundi Republic of Moldova Uruguay Gambia Mali Cabo Verde Marshall Islands Romania Uzbekistan Georgia Cambodia Ghana Mauritania Russian Federation Vanuatu Cameroon Greenland Mexico Rwanda Venezuela Central African Guam Micronesia (Federated Sao Tome and Principe (Bolivarian Republic Guatemala States of) Senegal Republic of) Chad Guinea Mongolia Sevchelles Viet Nam China Guinea-Bissau Sierra Leone Yemen Singapore Zambia

Source: World Health Organization, Global Health Observatory, Tuberculosis Incidence 2023. Countries with incidence rates of \geq 20 cases per 100,000 populations. For future updates, refer to http://www.who.int/tb/country/en/.



STUDENT HEALTH SERVICES

Zimbabwe

Imu.edu/health | Imuhealth@Imu.edu | (310) 338-2881 Burns Recreation Center, Ground-floor level, North