

# ENTRANCE REQUIREMENTS

ALL INCOMING STUDENTS **MUST** COMPLETE THE FOLLOWING BY December 1st 2023:

1. Review the Consent for Treatment Form.
2. Review the Notice of Privacy Practices.
3. Review the Telehealth Consent Form.
4. Provide documentation of two measles/mumps/rubella (MMR) immunization or proof of immunity (blood titer).
5. Complete Tuberculosis (TB) Screening and/or testing.

- **Go to MyHealth portal located at**

**(MyHealth.lmu.edu) login and click on the MEDICAL CLEARANCES**

*Tab located on the left hand side of your screen.*

- *Complete all items in your Medical Clearances section.*

**Please Note:**

- **Tuberculosis (TB) Screening and/or testing**

You must submit a TB skin or blood test done within the past 12 months, if your country of origin is designated by the World Health Organization (WHO) as a country with a high incidence of tuberculosis OR you answered "yes" to any of the questions listed in the TB Risk Screening Questionnaire in the Medical Clearances section.

- If your TB skin or blood test is positive, you must also submit a chest x-ray report as proof you do not have active TB.

*To view the list of countries designated by the WHO as being high risk for TB see reverse side of this document.*

## CONTACT STUDENT HEALTH SERVICES

**Phone:** 310.338.2881

**Fax:** 310.338.4417

**Email:**  
lmuhealth@lmu.edu

**Office:**

Burns Recreation Center  
Ground-floor level, North

**Mailing address:**

1 LMU Drive, MS 8455,  
Los Angeles CA 90045

### Questions?

Please review the  
Entrance Requirements  
FAQ's on our web page:  
[lmu.edu/health](http://lmu.edu/health)

*Allow 5 business days  
for SHS to process  
your forms.*



**Loyola Marymount  
University  
Student Health  
Services**

## HIGH RISK FACTORS FOR ACQUIRING TB ARE AS FOLLOW:

- Close contact with persons known or suspected to have active TB disease
- Persons born in one of the countries listed below that have a high incidence of active TB disease
- Frequent or prolonged visits to one or more of the countries listed below with a high prevalence of TB disease
- A resident and/or employee of high-risk congregate settings (e.g., correctional facilities, long-term care facilities, and homeless shelters)
- A volunteer or health-care worker who served clients who are at increased risk for active TB disease
- A member of any of the following groups that may have an increased incidence of latent M. tuberculosis infection or active TB disease — medically underserved, low-income, or abusing drugs or alcohol

### COUNTRIES WITH HIGH INCIDENCE OF TUBERCULOSIS PER WORLD HEALTH ORGANIZATION

Afghanistan	China, Hong Kong SAR	Guyana	Morocco	Solomon Islands
Algeria	China, Macao SAR	Haiti	Mozambique	Somalia
Angola	Colombia	Honduras	Myanmar	South Africa
Anguilla	Comoros	India	Namibia	South Sudan
Argentina	Congo	Indonesia	Nauru	Sri Lanka
Armenia	Cote d'Ivoire	Iraq	Nepal	Sudan
Azerbaijan	Democratic People's	Kazakhstan	Nicaragua	Suriname
Bangladesh	Republic of Korea	Kenya	Niger	Swaziland
Belarus	Democratic Republic	Kiribati	Nigeria	Tajikistan
Belize	of the Congo	Kuwait	Niue	Tanzania (United
Benin	Djibouti	Kyrgyzstan	Northern Mariana	Republic of)
Bhutan	Dominican Republic	Lao People's	Islands	Thailand
Bolivia(Plurinational)	Ecuador	Democratic Republic	Pakistan	Timor-Leste
State of)	El Salvador	Latvia	Palau	Togo
Bosnia and	Equatorial Guinea	Lesotho	Panama	Tokelau
Herzegovina	Eritrea	Liberia	Papua New Guinea	Trinidad and Tobago
Botswana	eSwatini	Libya	Paraguay	Tunisia
Brazil	Ethiopia	Lithuania	Peru	Turkmenistan
Brunei Darussalam	Fiji	Madagascar	Philippines	Tuvalu
Bulgaria	French-Polynesia	Malawi	Portugal	Uganda
Burkina Faso	Gabon	Malaysia	Qatar	Ukraine
Burundi	Gambia	Maldives	Republic of Korea	Uruguay
Cabo Verde	Georgia	Mali	Republic of Moldova	Uzbekistan
Cambodia	Ghana	Marshall Islands	Romania	Vanuatu
Cameroon	Greenland	Mauritania	Russian Federation	Venezuela
Central African	Guam	Mexico	Rwanda	(Bolivarian
Republic	Guatemala	Micronesia (Federated	Sao Tome and Principe	Republic of)
Chad	Guinea	States of)	Senegal	Viet Nam
China	Guinea-Bissau	Mongolia	Sierra Leone	Yemen
			Singapore	Zambia
				Zimbabwe

Source: World Health Organization, Global Health Observatory, Tuberculosis Incidence 2021. Countries with incidence rates of  $\geq 20$  cases per 100,000 populations. For future updates, refer to <http://www.who.int/tb/country/en/>.



**Loyola Marymount  
University**  
**Student Health  
Services**

### STUDENT HEALTH SERVICES

lmu.edu/health | [lmuhealth@lmu.edu](mailto:lmuhealth@lmu.edu) | (310) 338-2881  
Burns Recreation Center, Ground-floor level, North