

ENTRANCE REQUIREMENTS

ALL INCOMING STUDENTS **MUST** COMPLETE THE FOLLOWING BY JULY 15, 2020:

1. Review the Consent for Treatment Form.
2. Review the Notice of Privacy Practices.
3. Complete Tuberculosis (TB) Screening and/or testing.
4. Provide documentation of two measles/mumps/rubella (MMR) immunization or proof of immunity (blood titer).

Step 1: Go to MyHealth portal located at
(MyHealth.lmu.edu)

Step 2: Review and acknowledge the following forms

- **Consent for Treatment** and indicate agreement by checking the box. If you are under 18 years old, download **Consent for Treatment of a Minor** located in download forms. Print this form and have it signed by your parent or guardian and submit it to SHS.
- **Notice of Privacy Practices** and acknowledge by checking the box.

Step 3: TB testing

- If listed as required on MyHealth portal, complete the **TB Risk Screening Questionnaire**

Students born in a country with high incidence of TB (see list on the back) or students who answered "YES" to any of the questions outlined on the TB Risk Screening Questionnaire **Must:**

- Enter the date of TB skin or blood test done within the last 12 months on the **Immunization Entry form**.
- For positive blood or skin TB test you **MUST** submit a chest x-ray report.

Step 4: Complete Immunization Entry Form

- Input your immunization/testing dates.

Step 5: Upload Records

- Upload copies of your immunization/testing records and, if applicable, TB test records and chest x-ray report to **MyHealth** portal.

CONTACT STUDENT HEALTH SERVICES

Phone: 310.338.2881

Fax: 310.338.4417

Email:
lmuhealth@lmu.edu

Office:

Burns Recreation Center
Ground-floor level, North

Mailing address:

1 LMU Drive, MS 8455,
Los Angeles CA 90045

Questions?

Please review the
Entrance Requirements
FAQ's on our web page:
lmu.edu/health

*Allow 5 business days
for SHS to process
your forms.*



**Loyola Marymount
University
Student Health
Services**

HIGH RISK FACTORS FOR ACQUIRING TB ARE AS FOLLOW:

- Close contact with persons known or suspected to have active TB disease
- Persons born in one of the countries listed below that have a high incidence of active TB disease
- Frequent or prolonged visits to one or more of the countries listed below with a high prevalence of TB disease
- A resident and/or employee of high-risk congregate settings (e.g., correctional facilities, long-term care facilities, and homeless shelters)
- A volunteer or health-care worker who served clients who are at increased risk for active TB disease
- A member of any of the following groups that may have an increased incidence of latent M. tuberculosis infection or active TB disease — medically underserved, low-income, or abusing drugs or alcohol

COUNTRIES WITH HIGH INCIDENCE OF TUBERCULOSIS PER WORLD HEALTH ORGANIZATION

Afghanistan	China, Hong Kong SAR	Guyana	Morocco	Solomon Islands
Algeria	China, Macao SAR	Haiti	Mozambique	Somalia
Angola	Colombia	Honduras	Myanmar	South Africa
Anguilla	Comoros	India	Namibia	South Sudan
Argentina	Congo	Indonesia	Nauru	Sri Lanka
Armenia	Cote d'Ivoire	Iraq	Nepal	Sudan
Azerbaijan	Democratic People's	Kazakhstan	Nicaragua	Suriname
Bangladesh	Republic of Korea	Kenya	Niger	Swaziland
Belarus	Democratic Republic	Kiribati	Nigeria	Tajikistan
Belize	of the Congo	Kuwait	Niue	Tanzania (United
Benin	Djibouti	Kyrgyzstan	Northern Mariana	Republic of)
Bhutan	Dominican Republic	Lao People's	Islands	Thailand
Bolivia(Plurinational)	Ecuador	Democratic Republic	Pakistan	Timor-Leste
State of)	El Salvador	Latvia	Palau	Togo
Bosnia and	Equatorial Guinea	Lesotho	Panama	Tokelau
Herzegovina	Eritrea	Libya	Papua New Guinea	Trinidad and Tobago
Botswana	eSwatini	Lithuania	Paraguay	Tunisia
Brazil	Ethiopia	Madagascar	Peru	Turkmenistan
Brunei Darussalam	Fiji	Malawi	Philippines	Tuvalu
Bulgaria	French-Polynesia	Malaysia	Portugal	Uganda
Burkina Faso	Gabon	Maldives	Qatar	Ukraine
Burundi	Gambia	Mali	Republic of Korea	Uruguay
Cabo Verde	Georgia	Marshall Islands	Republic of Moldova	Uzbekistan
Cambodia	Ghana	Mauritania	Romania	Vanuatu
Cameroon	Greenland	Mexico	Russian Federation	Venezuela
Central African	Guam	Micronesia (Federated	Rwanda	(Bolivarian
Republic	Guatemala	States of)	Sao Tome and Principe	Republic of)
Chad	Guinea	Mongolia	Senegal	Viet Nam
China	Guinea-Bissau		Sierra Leone	Yemen
			Singapore	Zambia
				Zimbabwe

Source: World Health Organization, Global Health Observatory, Tuberculosis Incidence 2018. Countries with incidence rates of ≥ 20 cases per 100,000 populations. For future updates, refer to <http://www.who.int/tb/country/en/>.



**Loyola Marymount
University**
**Student Health
Services**

STUDENT HEALTH SERVICES

lmu.edu/health | lmuhealth@lmu.edu | (310) 338-2881
Burns Recreation Center, Ground-floor level, North

