

ENTRANCE REQUIREMENTS

ALL INCOMING STUDENTS **MUST** COMPLETE THE FOLLOWING BY July 1st 2022:

1. Review the Consent for Treatment Form.
2. Review the Notice of Privacy Practices.
3. Review the Telehealth Consent Form.
4. Provide documentation of two measles/mumps/rubella (MMR) immunization or proof of immunity (blood titer).
5. Complete Tuberculosis (TB) Screening and/or testing.

- **Go to MyHealth portal located at**

(MyHealth.lmu.edu) login and click on the MEDICAL CLEARANCES

Tab located on the left hand side of your screen.

- *Complete all items in your Medical Clearances section.*

Please Note:

- **Tuberculosis (TB) Screening and/or testing**

You must submit a TB skin or blood test done within the past 12 months, if your country of origin is designated by the World Health Organization (WHO) as a country with a high incidence of tuberculosis OR you answered "yes" to any of the questions listed in the TB Risk Screening Questionnaire in the Medical Clearances section.

- If your TB skin or blood test is positive, you must also submit a chest x-ray report as proof you do not have active TB.

To view the list of countries designated by the WHO as being high risk for TB see reverse side of this document.

CONTACT STUDENT HEALTH SERVICES

Phone: 310.338.2881

Fax: 310.338.4417

Email:
lmuhealth@lmu.edu

Office:

Burns Recreation Center
Ground-floor level, North

Mailing address:

1 LMU Drive, MS 8455,
Los Angeles CA 90045

Questions?

Please review the
Entrance Requirements
FAQ's on our web page:
lmu.edu/health

*Allow 5 business days
for SHS to process
your forms.*



**Loyola Marymount
University
Student Health
Services**

HIGH RISK FACTORS FOR ACQUIRING TB ARE AS FOLLOW:

- Close contact with persons known or suspected to have active TB disease
- Persons born in one of the countries listed below that have a high incidence of active TB disease
- Frequent or prolonged visits to one or more of the countries listed below with a high prevalence of TB disease
- A resident and/or employee of high-risk congregate settings (e.g., correctional facilities, long-term care facilities, and homeless shelters)
- A volunteer or health-care worker who served clients who are at increased risk for active TB disease
- A member of any of the following groups that may have an increased incidence of latent M. tuberculosis infection or active TB disease — medically underserved, low-income, or abusing drugs or alcohol

COUNTRIES WITH HIGH INCIDENCE OF TUBERCULOSIS PER WORLD HEALTH ORGANIZATION

<i>Afghanistan</i>	<i>China, Hong Kong SAR</i>	<i>Guyana</i>	<i>Morocco</i>	<i>Solomon Islands</i>
<i>Algeria</i>	<i>China, Macao SAR</i>	<i>Haiti</i>	<i>Mozambique</i>	<i>Somalia</i>
<i>Angola</i>	<i>Colombia</i>	<i>Honduras</i>	<i>Myanmar</i>	<i>South Africa</i>
<i>Anguilla</i>	<i>Comoros</i>	<i>India</i>	<i>Namibia</i>	<i>South Sudan</i>
<i>Argentina</i>	<i>Congo</i>	<i>Indonesia</i>	<i>Nauru</i>	<i>Sri Lanka</i>
<i>Armenia</i>	<i>Cote d'Ivoire</i>	<i>Iraq</i>	<i>Nepal</i>	<i>Sudan</i>
<i>Azerbaijan</i>	<i>Democratic People's</i>	<i>Kazakhstan</i>	<i>Nicaragua</i>	<i>Suriname</i>
<i>Bangladesh</i>	<i>Republic of Korea</i>	<i>Kenya</i>	<i>Niger</i>	<i>Swaziland</i>
<i>Belarus</i>	<i>Democratic Republic</i>	<i>Kiribati</i>	<i>Nigeria</i>	<i>Tajikistan</i>
<i>Belize</i>	<i>of the Congo</i>	<i>Kuwait</i>	<i>Niue</i>	<i>Tanzania (United</i>
<i>Benin</i>	<i>Djibouti</i>	<i>Kyrgyzstan</i>	<i>Northern Mariana</i>	<i>Republic of)</i>
<i>Bhutan</i>	<i>Dominican Republic</i>	<i>Lao People's</i>	<i>Islands</i>	<i>Thailand</i>
<i>Bolivia(Plurinational)</i>	<i>Ecuador</i>	<i>Democratic Republic</i>	<i>Pakistan</i>	<i>Timor-Leste</i>
<i>State of)</i>	<i>El Salvador</i>	<i>Latvia</i>	<i>Palau</i>	<i>Togo</i>
<i>Bosnia and</i>	<i>Equatorial Guinea</i>	<i>Lesotho</i>	<i>Panama</i>	<i>Tokelau</i>
<i>Herzegovina</i>	<i>Eritrea</i>	<i>Libya</i>	<i>Papua New Guinea</i>	<i>Trinidad and Tobago</i>
<i>Botswana</i>	<i>eSwatini</i>	<i>Lithuania</i>	<i>Paraguay</i>	<i>Tunisia</i>
<i>Brazil</i>	<i>Ethiopia</i>	<i>Madagascar</i>	<i>Peru</i>	<i>Turkmenistan</i>
<i>Brunei Darussalam</i>	<i>Fiji</i>	<i>Malawi</i>	<i>Philippines</i>	<i>Tuvalu</i>
<i>Bulgaria</i>	<i>French-Polynesia</i>	<i>Malaysia</i>	<i>Portugal</i>	<i>Uganda</i>
<i>Burkina Faso</i>	<i>Gabon</i>	<i>Maldives</i>	<i>Qatar</i>	<i>Ukraine</i>
<i>Burundi</i>	<i>Gambia</i>	<i>Mali</i>	<i>Republic of Korea</i>	<i>Uruguay</i>
<i>Cabo Verde</i>	<i>Georgia</i>	<i>Marshall Islands</i>	<i>Republic of Moldova</i>	<i>Uzbekistan</i>
<i>Cambodia</i>	<i>Ghana</i>	<i>Mauritania</i>	<i>Romania</i>	<i>Vanuatu</i>
<i>Cameroon</i>	<i>Greenland</i>	<i>Mexico</i>	<i>Russian Federation</i>	<i>Venezuela</i>
<i>Central African</i>	<i>Guam</i>	<i>Micronesia (Federated</i>	<i>Rwanda</i>	<i>(Bolivarian</i>
<i>Republic</i>	<i>Guatemala</i>	<i>States of)</i>	<i>Sao Tome and Principe</i>	<i>Republic of)</i>
<i>Chad</i>	<i>Guinea</i>	<i>Mongolia</i>	<i>Senegal</i>	<i>Viet Nam</i>
<i>China</i>	<i>Guinea-Bissau</i>		<i>Sierra Leone</i>	<i>Yemen</i>
			<i>Singapore</i>	<i>Zambia</i>
				<i>Zimbabwe</i>

Source: World Health Organization, Global Health Observatory, Tuberculosis Incidence 2020. Countries with incidence rates of ≥ 20 cases per 100,000 populations. For future updates, refer to <http://www.who.int/tb/country/en/>.



**Loyola Marymount
University**
**Student Health
Services**

STUDENT HEALTH SERVICES

lmu.edu/health | lmuhealth@lmu.edu | (310) 338-2881
Burns Recreation Center, Ground-floor level, North