

ENTRANCE REQUIREMENTS

ALL INCOMING STUDENTS **MUST** COMPLETE THE FOLLOWING BY January 7th, 2022:

1. Review the Consent for Treatment Form.
2. Review the Notice of Privacy Practices.
3. Complete Tuberculosis (TB) Screening and/or testing.
4. Provide documentation of two measles/mumps/rubella (MMR) immunization or proof of immunity (blood titer).

Step 1: Go to MyHealth portal located at
(MyHealth.lmu.edu)

Step 2: Review and acknowledge the following forms

- **Consent for Treatment** and indicate agreement by checking the box. If you are under 18 years old, download **Consent for Treatment of a Minor** located in download forms. Print this form and have it signed by your parent or guardian and submit it to SHS.
- **Notice of Privacy Practices** and acknowledge by checking the box.

Step 3: TB testing

- If listed as required on MyHealth portal, complete the **TB Risk Screening Questionnaire**

Students born in a country with high incidence of TB (see list on the back) or students who answered "YES" to any of the questions outlined on the TB Risk Screening Questionnaire **Must**:

- Enter the date of TB skin or blood test done within the last 12 months on the **Immunization Entry form**.
- For positive blood or skin TB test you **MUST** submit a chest x-ray report.

Step 4: Complete Immunization Entry Form

- Input your immunization/testing dates.

Step 5: Upload Records

- Upload copies of your immunization/testing records and, if applicable, TB test records and chest x-ray report to **MyHealth** portal.

CONTACT STUDENT HEALTH SERVICES

Phone: 310.338.2881

Fax: 310.338.4417

Email:
lmuhealth@lmu.edu

Office:

Burns Recreation Center
Ground-floor level, North

Mailing address:

1 LMU Drive, MS 8455,
Los Angeles CA 90045

Questions?

Please review the
Entrance Requirements
FAQ's on our web page:
lmu.edu/health

*Allow 5 business days
for SHS to process
your forms.*



Loyola Marymount
University
**Student Health
Services**

HIGH RISK FACTORS FOR ACQUIRING TB ARE AS FOLLOW:

- Close contact with persons known or suspected to have active TB disease
- Persons born in one of the countries listed below that have a high incidence of active TB disease
- Frequent or prolonged visits to one or more of the countries listed below with a high prevalence of TB disease
- A resident and/or employee of high-risk congregate settings (e.g., correctional facilities, long-term care facilities, and homeless shelters)
- A volunteer or health-care worker who served clients who are at increased risk for active TB disease
- A member of any of the following groups that may have an increased incidence of latent M. tuberculosis infection or active TB disease — medically underserved, low-income, or abusing drugs or alcohol

COUNTRIES WITH HIGH INCIDENCE OF TUBERCULOSIS PER WORLD HEALTH ORGANIZATION

<i>Afghanistan</i>	<i>China, Hong Kong SAR</i>	<i>Guyana</i>	<i>Morocco</i>	<i>Solomon Islands</i>
<i>Algeria</i>	<i>China, Macao SAR</i>	<i>Haiti</i>	<i>Mozambique</i>	<i>Somalia</i>
<i>Angola</i>	<i>Colombia</i>	<i>Honduras</i>	<i>Myanmar</i>	<i>South Africa</i>
<i>Anguilla</i>	<i>Comoros</i>	<i>India</i>	<i>Namibia</i>	<i>South Sudan</i>
<i>Argentina</i>	<i>Congo</i>	<i>Indonesia</i>	<i>Nauru</i>	<i>Sri Lanka</i>
<i>Armenia</i>	<i>Cote d'Ivoire</i>	<i>Iraq</i>	<i>Nepal</i>	<i>Sudan</i>
<i>Azerbaijan</i>	<i>Democratic People's</i>	<i>Kazakhstan</i>	<i>Nicaragua</i>	<i>Suriname</i>
<i>Bangladesh</i>	<i>Republic of Korea</i>	<i>Kenya</i>	<i>Niger</i>	<i>Swaziland</i>
<i>Belarus</i>	<i>Democratic Republic</i>	<i>Kiribati</i>	<i>Nigeria</i>	<i>Tajikistan</i>
<i>Belize</i>	<i>of the Congo</i>	<i>Kuwait</i>	<i>Niue</i>	<i>Tanzania (United</i>
<i>Benin</i>	<i>Djibouti</i>	<i>Kyrgyzstan</i>	<i>Northern Mariana</i>	<i>Republic of)</i>
<i>Bhutan</i>	<i>Dominican Republic</i>	<i>Lao People's</i>	<i>Islands</i>	<i>Thailand</i>
<i>Bolivia(Plurinational)</i>	<i>Ecuador</i>	<i>Democratic Republic</i>	<i>Pakistan</i>	<i>Timor-Leste</i>
<i>State of)</i>	<i>El Salvador</i>	<i>Latvia</i>	<i>Palau</i>	<i>Togo</i>
<i>Bosnia and</i>	<i>Equatorial Guinea</i>	<i>Lesotho</i>	<i>Panama</i>	<i>Tokelau</i>
<i>Herzegovina</i>	<i>Eritrea</i>	<i>Libya</i>	<i>Papua New Guinea</i>	<i>Trinidad and Tobago</i>
<i>Botswana</i>	<i>eSwatini</i>	<i>Lithuania</i>	<i>Paraguay</i>	<i>Tunisia</i>
<i>Brazil</i>	<i>Ethiopia</i>	<i>Madagascar</i>	<i>Peru</i>	<i>Turkmenistan</i>
<i>Brunei Darussalam</i>	<i>Fiji</i>	<i>Malawi</i>	<i>Philippines</i>	<i>Tuvalu</i>
<i>Bulgaria</i>	<i>French-Polynesia</i>	<i>Malaysia</i>	<i>Portugal</i>	<i>Uganda</i>
<i>Burkina Faso</i>	<i>Gabon</i>	<i>Maldives</i>	<i>Qatar</i>	<i>Ukraine</i>
<i>Burundi</i>	<i>Gambia</i>	<i>Mali</i>	<i>Republic of Korea</i>	<i>Uruguay</i>
<i>Cabo Verde</i>	<i>Georgia</i>	<i>Marshall Islands</i>	<i>Republic of Moldova</i>	<i>Uzbekistan</i>
<i>Cambodia</i>	<i>Ghana</i>	<i>Mauritania</i>	<i>Romania</i>	<i>Vanuatu</i>
<i>Cameroon</i>	<i>Greenland</i>	<i>Mexico</i>	<i>Russian Federation</i>	<i>Venezuela</i>
<i>Central African</i>	<i>Guam</i>	<i>Micronesia (Federated</i>	<i>Rwanda</i>	<i>(Bolivarian</i>
<i>Republic</i>	<i>Guatemala</i>	<i>States of)</i>	<i>Sao Tome and Principe</i>	<i>Republic of)</i>
<i>Chad</i>	<i>Guinea</i>	<i>Mongolia</i>	<i>Senegal</i>	<i>Viet Nam</i>
<i>China</i>	<i>Guinea-Bissau</i>		<i>Sierra Leone</i>	<i>Yemen</i>
			<i>Singapore</i>	<i>Zambia</i>
				<i>Zimbabwe</i>

Source: World Health Organization, Global Health Observatory, Tuberculosis Incidence 2018. Countries with incidence rates of ≥ 20 cases per 100,000 populations. For future updates, refer to <http://www.who.int/tb/country/en/>.



**Loyola Marymount
University**
**Student Health
Services**

STUDENT HEALTH SERVICES

lmu.edu/health | lmuhealth@lmu.edu | (310) 338-2881
Burns Recreation Center, Ground-floor level, North

